

Amendments to the Specification:

Please replace the paragraph at page 13, line 17 to page 14, line 2, with the following amended paragraph:

The magnitude of a prophylactic or therapeutic dose of the active ingredient (e.g., S-tofisopam or salt thereof) or S-tofisopam prodrug and, if desired, other anticonvulsant for treating or preventing convulsions or seizures will vary with the severity of the patient's affliction and the route of administration. The dose and dose frequency will also vary according to the age, weight and response of the individual patient. In general, the recommended daily dose range for the conditions described herein can lie within the range of from approximately 10 mg to approximately 1200 mg per day, generally divided equally into doses given one to four times a day. A daily dose range can be between 50 mg and 600 mg per day, usually divided equally into a one to four times a day dosing. Alternatively, a daily dose range can be between 100 mg and 400 mg per day, usually divided equally into a two to four times a day dosing. Also, when based on the body weight of the patient, S-tofisopam may be administered at a dosage of less than 30 mg/kg of the patient's body weight. It can be necessary to use dosages outside these ranges in some cases and adjust the amounts of S-tofisopam, salt or prodrug thereof administered alone or in combination with the other anti-convulsant(s). The treating physician will know how to increase, decrease or interrupt treatment based upon patient response. The various terms described above such as "therapeutically effective amount," are encompassed by the above-described dosage amounts and dose frequency schedule.